

## Weight Decrease in Full-Term Newborns in the First 48 Postnatal Hours

Miguel Gallardo López<sup>1\*</sup>, Esteban Gallardo Cadenasso<sup>2</sup> and Laura Gallardo Cadenasso<sup>3</sup>

<sup>1</sup>Neonatologist, Resident at the Neonatology Unit, Clínica Dávila, Santiago, Chile

<sup>2</sup>Public Policy Master's Student, Universidad de Chile, Santiago, Chile

<sup>3</sup>Nursing Student, Universidad del Desarrollo, Santiago, Chile

**\*Corresponding Author:** Miguel Gallardo López, Neonatologist, Resident at the Neonatology Unit, Clínica Dávila, Santiago, Chile.

**Received:** May 28, 2018; **Published:** September 28, 2018

### Abstract

**Introduction:** It is known that the newborn (NB) presents weight loss immediately after birth. The magnitude of the weight loss is a controversial subject and there is no data in our environment.

**Objectives:** Determining the percentage of daily weight loss in healthy full-term newborns in the first 48 hours of being born. Specifying whether the factors: gender, type of delivery, adequacy to gestational age and use of milk formula influences weight loss.

**Patients and Method:** A prospective descriptive study in healthy full-term newborns with  $\geq 37$  weeks of gestational age during their stay in nursery. Weights were collected at birth, 24 and 48 hours. After this, the percentages of weight loss were calculated.

**Results:** 2960 NB were analyzed. The average weight loss on the first day was 4.43%, the second day was of 2.51% and the total decline at 48 hours was of 6.85%. The determining factor in weight loss was the type of delivery. The NB delivered by cesarean showed a greater weight loss in the first 24 to 48 hours. The use of milk formula determines a smaller weight loss in the NB compared to those fed with breast milk (BM).

**Conclusions:** The percentage of weight loss per day of the NB in our center is alike to that stated in international literature. A cesarean delivery is the factor associated with greater weight loss. The use of milk formula determines lower decrease of weight.

**Keywords:** Postnatal Weight Loss; Exclusive Breastfeeding; Neonates; Jaundice

### Introduction

Most healthy newborns show a period of weight loss right after birth [1]. This weight loss during the first days of life is something known; its cause and magnitude are subject to controversy [1,2]. It is suggested that on average children lose 4% - 7% of their birth weight [3] and that they recover it around the third day of life [4].

Breastfed newborns are only fed with small amounts of liquids during the first days of life [5,6], therefore they tend to lose weight before the weight increase occurs [7,8]. Excessive weight loss can be a sign of a low milk supply or of insufficient milk transference [9,10]. The most common way to evaluate if breastfeeding is effective and efficient is to weigh the newborn during its stay in the nursery (post-partum period) [11,12]. Data based on the evidence related to the weight loss of the newborn during the first days of life are limited [10].

The Mother-and-Baby-Friendly Initiative, which was enacted by the OMS in cooperation with UNICEF, recommends exclusive breastfeeding during the hospitalization of the newborn and that no formula or water should be added [13,14].

In breastfed newborns, the enteral intake is low during the production period of colostrum, and therefore the initial weight loss after birth is almost universal [5,6,15]. Even though this short period of low enteral intake and weight loss is well tolerated by most newborns, some of them develop complications due to weight loss, such as hyperbilirubinemia and dehydration [16-19].

It is essential to identify what is considered as normal weight loss in the newborn since it will give doctors more information, which can help to make a well-grounded decision of adding additional milk formula to breastfeeding.

### Objectives of the Study

The main objectives of this study are: to determine the percentage of daily weight loss of healthy newborns during their stay in the nursery (postpartum period) and to specify if gender, type of birth, gestational age, adequacy for gestational age, and use of milk formula affect the loss of weight.

### Patient and Method

**Design:** Prospective cohort study. Data gathered from two periods were analyzed: from November 2010 to January 2011 and from January 2014 to March 2014. The following newborns were excluded: term newborns younger than 37 weeks of gestational age and newborns who stayed in the nursery with their mother for at least 48 hours. All patients were born in the maternity unit of Clínica Dávila, Santiago. Two different periods of time were analyzed in order to increase the size of the sample and to avoid sampling bias. There were no variations in the management of the patients during both periods.

Nursery, or postpartum period, is the clinical instance where newborns are cared for and monitored, from their birth to their discharge. During this period, the patient stays with their mother in a single or shared room for 24 hours. He or she is also under monitoring by midwives and paramedic technicians trained in couplet care.

The following newborns were excluded from the analysis: all preterm newborns, multiple pregnancy, and patients that required hospitalization in the neonatology unit.

**Clinical data:** Birth weights were recorded at 24 and 48 hours after birth. All weights were measured with the same digital weight scale, Seca 334, which was calibrated daily as standard of service. Data were gathered every day from the daily nursing reports. In addition, the following data were recorded: gender, type of birth, birth adequacy, and type of feeding (exclusive breastfeeding, artificial formula or mixed feeding).

Regarding the use and indication of artificial formula, there are no policies or protocols on the use of supplements in our Clinic and the indication of artificial feeding was made according to individual medical criteria without influences, from the researchers, on the decision.

**Ethical aspects:** The Scientific Ethics Committee became aware of the study and approved its development in Clínica Dávila and allowed the registration of data from the patients and the subsequent publication of the results.

**Statistical analysis:** The data were analyzed with STATA 13.0 software. In order to standardize measurements, weights were recorded in grams and then the percentage of weight loss in relation to birth weight and last day weight was calculated. The maximum percentage of daily weight loss was analyzed. A multivariate regression model of the evolution of weight loss percentages was performed.

### Results

The total sample was of 2,960 newborns, 1,149 from the 2010 - 2011 period and 1,822 from the 2014 period. The average gestational age was 38.78 weeks, ranging from 37 to 41 weeks (SD: 0.954). The type of birth showed that 51.93% are cesarean, 40.27% are normal delivery and 7.81% are assisted delivery (forceps). The distribution according to gender was 50.74% males (Table 1).

Regarding adequacy, 83.18% were adequate for gestational age (AGA), 10.51% were large for gestational age (LGA) and 6.3% were small for gestational age (SGA) (Table 1). None of the previously described variables showed differences in both periods (gestational age, type of birth, gender and adequacy for gestational age).

From the sample, 1,151 (39.39%) newborns received exclusive breastfeeding (EBF) during their stay in the nursery, 1,809 (61.66%) newborns received formula one or more times (Table 1). A higher percentage of newborns received EBF in the first period than in the second period (54.05% and 29.27%, respectively), which is statistically relevant.

Variables		Nº of patients	Percentage
Gestational Age (Weeks)	37	248	8,38
	38	916	30,95
	39	1058	36,00
	40	679	22,94
	41	59	1,99
Type of Delivery	Vaginal	1192	40,27
	Cesarean	1537	51,93
	Spatula	147	4,97
	Forceps	84	2,84
Gender	Female	1458	49,26
	Male	1502	50,74
Adecuacy	Adequate	2462	83,18
	Large	311	10,51
	Small	187	6,32
Type of Feeding	Breast	1151	38,89
	Formula	1809	61,11
	Total	2960	100,00

**Table 1:** General Characteristics of the 2960 FTN.

*FTN indicates full-term newborns.*

The weight loss analysis of the studied sample showed that during the first 24 hours the average weight loss was 4.43% (SD: 1.96), the weight loss on the second day was 2.51% (SD: 1.86) compared to the first day and the total weight loss during the first 48 hours was 6.85% (SD: 1.92) compared to the birth weight (Figure 1). By analyzing each period, the results were similar: in the 2010 - 2011 period, the weight loss during the first 24 hours was 4.4% (SD: 2.03), the weight loss on the second day was 2.49% (SD: 1.96) compared to the first day and the total weight loss during the first 48 hours was 6.85% (SD: 1.95) compared to the birth weight, while in the 2014 period, 4.42% (SD: 1.92) weight loss during the first 24 hours, 2.52% (SD: 1.79) weight loss on the second day and a 6.86% (SD: 1.9) total weight loss during the first 48 hours, compared to the birth weight.



Figure 1: Percentage of Weight Loss on Postnatal Days One and Two.

After analyzing which studied factors affect weight loss, the multivariate study of the first 24 hours shows that there was no significant difference in any of the studied variables (Table 2).

The analysis of the period between the 24 and 48 hours of life shows that newborns who are fed with artificial formula lose more weight than newborns fed with EBF (P = 0.033); newborns delivered by cesarean lose more weight than newborns delivered by normal delivery (P = 0.000). Male newborns lose less weight than female newborns (P = 0.000). SGA newborns lose less weight than AGA newborns (P = 0.000). Gestational age also shows a statistical difference, the higher the gestational age, the lower the weight loss (P = 0.006) (Table 2).

The total weight loss analysis of the first 48 hours of life shows that there were no statistical differences in gender, gestational age, and adequacy for gestational age. However, there were differences in the type of birth and use of artificial formula, since newborns delivered by cesarean lose more weight than newborns delivered by normal delivery (P= 0.039) and newborns fed with formula lose less weight than newborns fed with EBF (P = 0.039) (Table 2).

	Day 1 Coefficient	P	Day 2 Coefficient	P	Total Coefficient	P
Mixed Feeding	466,712	p < 0,1	-0,1520745	p < 0,05	0,2482842	p < 0,05
Cesarean	673,5744	p < 0,1	-0,7251474	p < 0,01	-0,2004924	p < 0,05
Males	667,7991	p < 0,1	0,2465522	p < 0,01	0,0247866	p < 0,1
Adequacy						
SGA	-337,0152	p < 0,1	0,8487974	p < 0,01	0,3432167	p < 0,1
LGA	-612,9448	p < 0,1	-0,0520678	p < 0,1	-0,0074705	p < 0,1
> Gestacional Age	484,1344	p < 0,1	0,099387	p < 0,01	0,0285867	p < 0,1

**Table 2:** Factors Analyzed in Weight Loss.

*SGA indicates small for gestational age*

*LGA indicated large for gestational age*

### Discussion

This study shows that healthy newborns experience a 6.85% average weight loss in the first 48 hours of life, which is similar to what is reported in international literature [2,3,10,15,26]. The first studies that reported this phenomenon were made by Masiels., *et al.* [20] in 1980, where they reported a 6% average weight loss in the first three days of life [20] and in a second study, they reported a 6.86% average weight loss, but did not indicate if it happened in two or three days [3]. Furthermore, these findings came from studies performed in newborns that were made to evaluate neonatal jaundice and risk factors which influence their development; therefore, there is bias regarding the chosen population and those studies do not have the required amount of patients to estimate precisely the normal limits of weight loss during the first days of life. In later studies performed by Chantry., *et al.* [21], they report a 5.8% ± 3.7% loss on the third day [21].

The percentage of weight loss analysis provides the following data: the highest percentage of weight loss occurs on the first day of life, an average of 4.43%; a lower weight loss occurs on the second day, an average of 2.81% compared to the first day. A systematic review by Noel-Weiss., *et al.* [10] which included 11 studies performed on healthy term newborns, fed with exclusive breastfeeding, determined that the average weight loss varies from 5.7% to 6.6%, with a 2% standard deviation, the higher percentages of weight loss appeared on the second and third day after birth; the studies incorporated in this review are difficult to compare, since there are different statistical descriptions to express the weight changes [10].

After analyzing the risk factors that affect weight loss, we found that the type of birth is a decisive factor in the weight loss of newborns. Delivery by cesarean was a decisive factor in higher weight loss between the 24 and 48 hours of life and in the total weight loss during the first 48 hours of life. These findings coincide with what was reported by Maganaro., *et al.* [19], where the association between delivery by cesarean and a higher weight loss was described [19]. Delivery by cesarean has been described as a risk factor for suboptimal breastfeeding, a delay for the start of breastfeeding [22-24] and excess weight loss [19]. Regnault., *et al.* [25] showed that a newborn delivered by cesarean was 2.42 more likely to have excess weight loss [25]. These findings indicate that pre-birth factors, such as dehydration, can affect the postnatal weight loss of breastfed newborns [21,26]. The higher weight loss is associated with an intravenous fluid overload in mothers undergoing cesarean delivery with the subsequent over-hydration of NB, which contributes to increased weight loss [27]. Chantry., *et al.* [21] reported that the relative risk of newborns who lost > 10% of their birth weight increased three times when women had a balance of more than a 200 mL/h positive liquid during the intrapartum period [21]. Watson., *et al.* [28] suggested the existence of intravenous liquids threshold received by the mother during birth, which affects the newborn weight loss. Therefore, the volume of intravenous liquids is a factor which contributes to the early weight loss of newborns during the first 48 hours of life [28].

Regarding the use of formula in our study, after analyzing the period between 24 and 48 hours, it was observed that newborns fed with formula had a higher weight loss than newborns fed with EBF [2,15,27]. Maisels, *et al.* [3] determined that newborns fed with breastfeeding lost a  $6.86\% \pm 2.97\%$  average of their birth weight, while newborns fed with formula lost  $4.24 \pm 2.88\%$  ( $P < 0,02$ ). Macdonald, *et al.* [15] demonstrated that breastfed newborns lose more weight and recover their weight slower than newborns fed with formula. The results analyzed regarding the use of formula may seem contradictory when comparing both periods, we found that newborns fed with formula had a higher weight loss than breastfed newborns after analyzing the 24 - 48 hours. This could be explained by an inverse causality since newborns often lose an excess of weight, the first approach is to supplement breastfeeding with formula, and therefore, the use of formula will be a consequence of the weight loss and not its cause. On the other hand, Fonseca, *et al.* [2] determined that breastfed newborns did not lose weight, which indicates that breast milk is enough to support the growth of the newborns during the first hours of life; in contrast, newborns fed with formula or mixed feeding were more likely to have a sub-optimal weight loss, which could indicate overfeeding [2].

After analyzing the risk factors in the 24 - 48 hour period, it was observed that male SGA and GAG newborns had a lower weight loss than female AGA and SGA newborns. Marten, *et al.* [27] stated that birth weight may be an important factor in weight loss in the immediate neonatal period, presenting a positive association with the weight loss [25]. Fonseca, *et al.* [2] determined that an insufficient weight loss is associated with birth weight, newborns who weigh  $< 2,500\text{g}$  were 2.68 times more likely to lose weight [2]. This can be due to a recovery of growth, which could produce an increased growth in comparison with the normal rates for that age after a period of conditions that inhibit growth, which is common in SGA newborns [29]. Chapman and Pérez-Escamilla [22] found a higher risk of delayed breastfeeding in mothers whose newborn weighed less than 3.6 kg (8 lb) [22].

After analyzing the studied risk factors and evaluating in which way they affect weight loss, the multivariate analysis of the first 24 hours shows that there was no significant difference in any of the analyzed variables. This can be explained by the absence of some variables in our study, such as labor time, the amount of administrated fluids and anesthesia doses, which have been demonstrated in literature to affect weight loss [9,21,28] and can be especially considered risk factors of weight loss in newborns during the first 24 hours of life.

This study has the limitation of not having considered variables related to the mother, such as parity, previous breastfeeding experience, labor time, type of anesthesia, which have been described as risk factors associated to higher weight loss [19,26]. Another limitation is that it did not consider the use of fluids during the intrapartum period, where the administration would cause an expansion of the fetal volume and determine a higher weight loss [21].

The strength of this study is the sample size. The birth weight was recorded as a habitual procedure in the clinical practice and all weight measurements were performed prospectively, which reduces recall bias.

A protocol that addresses and evaluates weight loss during the first 48 hours of life should always take into consideration factors that in our study lead to a higher risk of presenting high weight loss, such as the type of birth, gender, gestational age and adequacy for gestational age [30].

### Conclusion

It was possible to determine the percentage of daily weight loss of newborns in our center. The percentage of daily weight loss of newborns in our center was similar to what has been reported in international literature. Cesareans are a factor which are associated with higher weight loss. The use of milk formula determines lower weight loss in newborns.

The results of this study strengthen the need for attention and special follow-ups of couplets with the risk of presenting excess weight loss after birth.

We believe that the appropriate use of our data will help to substantiate interventions, such as supplementing formula.

### Acknowledgments

Manuela Hume Cadenasso, English-Spanish Translation Student, Universidad Chileno-Británica de Cultura, Santiago, Chile.

### Bibliography

1. Wright CM and Parkinson KN. "Postnatal weight loss in term infants: ¿what is normal and do growth charts allow for it?" *Archives of Disease in Childhood. Fetal and Neonatal Edition* 89.3 (2004): F254-F257.
2. Fonseca MJ., *et al.* "Determinants of weight changes during the first 96 hours of life in full-term newborns". *Birth* 41.2 (2014): 160-168.
3. Maisels MJ., *et al.* "Jaundice in the healthy newborn infant: a new approach to an old problem". *Pediatrics* 81.4 (1988): 505-511.
4. Marchini G and Stock S. "Thirst and vasopressin secretion counteract dehydration in newborn infants". *Journal of Pediatrics* 130.5 (1997): 736-739.
5. Slusher TM., *et al.* "Comparison of maternal milk (breastmilk) expression methods in an African nursery". *Breastfeeding Medicine* 7.2 (2012): 107-111.
6. Saint L., *et al.* "The yield and nutrient content of colostrum and milk of women from giving birth to 1 month post-partum". *British Journal of Nutrition* 52.1 (1984): 87-95.
7. Neville MC and Morton J. "Physiology and endocrine changes underlying human lactogenesis II". *Journal of Nutrition* 131.11 (2001): 3005S-3008S.
8. Hartmann PE. "Lactation and reproduction in Western Australian women". *Journal of Reproductive Medicine* 32.7 (1987): 543-547.
9. Dewey KG., *et al.* "Delayed lactogenesis and excess neonatal weight loss are common across ethnic and socioeconomic categories of primiparous women in northern California [abstract]". *The FASEB Journal* 23.1 (2009): 344.
10. Noel-Weiss J., *et al.* "Physiological weight loss in the breastfed neonate: a systematic review". *Open Medicine* 2.4 (2008): e99-e110.
11. Harding D., *et al.* "Hypernatraemia: why bother weighing breast fed babies?" *Archives of Disease in Childhood. Fetal and Neonatal Edition* 85.2 (2001): F145.
12. Oddie S., *et al.* "Hypernatraemic dehydration and breast feeding: a population study". *Archives of Disease in Childhood* 85.4 (2001): 318-320.
13. UNICEF/WHO. "Baby-Friendly Hospital Initiative: Revised, Updated and Expanded for Integrated Care, Section 1, Background and Implementation, Preliminary version" (2006).
14. "American Academy of Pediatrics Section on Breastfeeding. Breastfeeding and the use of human milk". *Pediatrics* 129.3 (2012): e827-e841.
15. Macdonald PD., *et al.* "Neonatal weight loss in breast and formula fed infants". *Archives of Disease in Childhood. Fetal and Neonatal Edition* 88.6 (2003): F472-F476.
16. van Dommelen P., *et al.* "Reference chart for relative weight change to detect hypernatraemic dehydration". *Archives of Disease in Childhood* 92.6 (2007): 490-494.

17. Paul IM., *et al.* "Preventable newborn readmissions since passage of the Newborns' and Mothers' Health Protection Act". *Pediatrics* 118.6 (2006): 2349-2358.
18. Chen HL., *et al.* "Neonatal readmission within 2 weeks after birth". *Acta Paediatrica Taiwanica* 46.5 (2005): 289-293.
19. Manganaro R., *et al.* "Incidence of dehydration and hypernatremia in exclusively breast-fed infants". *Journal of Pediatrics* 139.5 (2001): 673-675.
20. Maisels MJ and Gifford K. "Breast-feeding, weight loss and jaundice". *Journal of Pediatrics* 102.1 (1983): 117-118.
21. Chantry C., *et al.* "Excess weight loss in first-born breastfed newborns relates to maternal intrapartum fluid balance". *Pediatrics* 127.1 (2011): e171-e179.
22. Chapman DJ and Pérez-Escamilla R. "Identification of risk factors for delayed onset of lactation". *Journal of The American Dietetic Association* 99.4 (1999): 450-454.
23. Hildebrandt HM. "Maternal perception of lactogenesis time: a clinical report". *Journal of Human Lactation* 15.4 (1999): 317-323.
24. Vestermark V., *et al.* "Influence of the mode of delivery on initiation of breast-feeding". *European Journal of Obstetrics and Gynecology and Reproductive Biology* 38.1 (1991): 33-38.
25. Regnault N., *et al.* "Determinants of neonatal weight loss in term-infants: specific association with pre-pregnancy maternal body mass index and infant feeding mode". *Archives of Disease in Childhood. Fetal and Neonatal Edition* 96.3 (2011): F217-F222.
26. Dewey K., *et al.* "Risk factors for suboptimal infant breastfeeding behavior, delayed onset of lactation, and excess neonatal weight loss". *Pediatrics* 112 (2003): 607-619.
27. Martens PJ and Romphf L. "Factors associated with newborn in hospital weight loss: Comparisons by feeding method, demographics, and birthing procedures". *Journal of Human Lactation* 23.3 (2007): 233-245.
28. Watson J., *et al.* "A randomized controlled trial of the effect of intrapartum intravenous fluid management on breastfed newborn weight loss". *Journal of Obstetric, Gynecologic, and Neonatal Nursing* 41.1 (2012): 24-32.
29. Gafni RI and Baron J. "Catch-up growth: Possible mechanisms". *Pediatric Nephrology* 14.7 (2000): 616-619.
30. Flaherman VJ., *et al.* "Early weight loss nomograms for exclusively breastfed newborns". *Pediatrics* 135.1 (2015): e16-e23.

**Volume 7 Issue 10 October 2018**

**©All rights reserved by Miguel Gallardo López., *et al.***